

**WARNING**

**KEEP OUT OF REACH OF CHILDREN**

**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

mondo®

A stylized graphic of a plant with three leaves. The top leaf is dark green, and the two bottom leaves are a lighter green. The leaves are positioned behind the 'mondo' text, with the top leaf partially overlapping the 'o'.

# FUNGICIDE

**Contains:** 375 g/litre trifloxystrobin plus 160 g/litre cyproconazole in the form of a suspension concentrate

**For disease control in fodder and sugar beets**

NET CONTENTS: 1L, 5L, 10L, 20L

©Registered Trademark of Arxada NZ Limited

Approved pursuant to the HSNO Act 1996, No: HSR101053. See [www.epa.govt.nz](http://www.epa.govt.nz) for approval conditions

Registered pursuant to the ACVM Act 1997, No: P009303. See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions

**arxada**

	GHS CLASSIFICATION	HAZARD STATEMENT
	Acute oral toxicity Category 4, Skin sensitisation Category 1, Reproductive toxicity Cat. 2  Specific target organ toxicity (repeated exposure) Category 2,  Hazardous to terrestrial vertebrates, Hazardous to the aquatic environment acute & chronic Cat. 1,	Harmful if swallowed. May cause an allergic skin reaction. Suspected of damaging fertility or the unborn child. May cause damage to organs through prolonged or repeated exposure  Hazardous to terrestrial vertebrates. Very toxic to aquatic life, with long lasting effects.
<p><b>FIRST AID:</b> IF EXPOSED OR CONCERNED: Call the National Poisons Centre 0800 POISON (0800 764 766) / doctor / physician / first aider.            IF SWALLOWED: Rinse Mouth. Call the National Poisons Centre / doctor / physician / first aider. Seek medical help if you feel unwell.            IF ON SKIN: Wash with plenty of water. If skin irritation or rash occurs: Get medical advice/attention.            Take off contaminated clothing and wash it before reuse.</p> <p><b>PERSONAL PROTECTION:</b> Do not breathe mist/vapours/spray. Wear protective gloves/protective clothing/eye protection/face protection. Do not eat, drink or smoke while using this product. Contaminated work clothing should not be allowed out of the workplace. Wash all exposed external body areas thoroughly after handling.</p> <p><b>STORAGE:</b> Store locked up. Store in original container, tightly closed and in a dry, cool area away from foodstuffs. Store in accordance with NZS 8409 Management of Agrichemicals.</p> <p><b>DISPOSAL:</b> Avoid contamination of any water supply with product or empty container. Triple rinse empty container placing rinse water in the spray tank. If recycling, discard cap and deliver clean container to an Agrecovery depot or crush and bury in an approved landfill. Dispose of product only by using according to this label, or at an approved landfill or disposal facility.</p> <p><b>SPILLAGE:</b> Wear appropriate protective clothing. Exclude all bystanders from the vicinity of the spill and prevent material from entering waterways. Contain liquid spills and absorb with inert material and place in waste containers. Wash area with water and absorb with further inert material. Dispose of waste safely in an approved landfill.</p> <p><b>SAFETY DATA SHEET:</b> Additional information is listed in the Safety Data Sheet available from Arxada NZ Limited or <a href="http://www.arxada.co.nz">www.arxada.co.nz</a></p> <p><b>QUALIFICATION REQUIREMENTS:</b> Persons mixing, loading, applying, or otherwise handling Mondo must meet qualification requirements set out in the EPA Hazardous Substances (Hazardous Property Controls) Notice 2017.</p>		

**PLEASE READ ENTIRE LABEL BEFORE MIXING OR APPLYING. SHAKE OR STIR WELL BEFORE USE.**

**GENERAL INFORMATION:** Mondo combines two effective fungicides that both offer protectant and systemic activity for use as a preventative treatment for disease control in fodder beet and sugar beet.

**It is an offence to use this product on animals.**

**DIRECTIONS FOR USE**

CROP	DISEASE	RATE/ha	CRITICAL COMMENTS
FODDER BEET SUGAR BEET	Rust Powdery mildew Cercospora Leaf Spot Ramularia Leaf Spot	350 ml	Make the first application as soon as the disease becomes active in the crop, conditions are likely to be favourable for its spread and before it becomes well established. A second application may be applied no sooner than 21 days later if these disease conditions are repeated.

**MIXING:** Always shake container or stir contents well before use. Slowly pour the correct amount of Mondo into the partly filled tank, and while agitating add the remainder of the water. Add any tank mixture products separately and adjuvant last. Maintain agitation during spraying and any stoppages. Only mix sufficient spray for immediate use. DO NOT leave the spray mix in the sprayer overnight.

**COMPATIBILITY:** Mondo is compatible with the insecticide Cyhella encaps® and the foliar fertiliser Boron 150 in tank mixture. Mondo is compatible with Beatrix® but seek advice from herbicide manufacturers for any other herbicide mixtures. Refer to the tank mixture product labels for further instructions. Each product should be added separately to the bulk of the water in the spray tank while filling with the agitator working.

**ADJUVANTS:** Always add a suitable adjuvant, such as a non-ionic wetter at label rates.

**APPLICATION:** Mondo can be applied through all conventional ground rig and aerial spray equipment correctly calibrated to achieve good coverage of target surfaces. The chosen spray volume and the sprayer set up and operation may need to be changed as the crop grows and matures. It is essential that good coverage of the total leaf canopy is achieved. Use at least 200 litres/ha for ground rigs and at least 80 litres/ha via aerial methods. Do not spray over or onto any water bodies.

**RAINFALL:** Effectiveness may be reduced if significant rainfall occurs within three hours of application.

**It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.**

**WITHHOLDING PERIOD:**

Fodder beet: Six weeks

**RESISTANCE MANAGEMENT:** **GROUP 11 + 3 FUNGICIDE**

Mondo contains trifloxystrobin, a QoI (quinone outside inhibitor) mode of action group 11 fungicide plus cyproconazole, a DMI (demethylation inhibitor) mode of action group 3 fungicide. Despite having two different modes of action certain strains of disease causing organisms may become tolerant to this fungicide due to natural variation and selection. If Mondo is used exclusively these strains may dominate a population and cause fungicide failure. To minimise any possible resistance risk use strictly in accordance with the label instructions. Apply Mondo preventatively or as early as possible in the disease cycle and use fungicides with a different mode of action at other times. For further information see: [www.resistance.nzpps.org](http://www.resistance.nzpps.org)

**NOTICE TO BUYER:** The manufacturer and vendors of Mondo warrant the formulation to contain the specified active ingredients within accepted tolerance but make no warranty of any kind on the uses of material and accept no responsibility, whether the material is used in accordance with the label directions or not.

**REGISTERED TO:** Arxada NZ Limited, Hudson Road,  
 New Plymouth, NZ.  
 Tel: 06-755-9234 Fax: 06-755-1174  
**EMERGENCY PHONE: 0800 CHEMCALL 0800 243 622**

D.O.M:  
 BATCH NO:



Mondo 221017