

**DANGER**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE**  
**OPENING OR USING**



# Jive®

221122

## HERBICIDE

**Contains:** 70.5 g/litre fenoxaprop-P-ethyl in the form of an emulsifiable concentrate  
Also contains: 35 g/litre cloquintocet-mexyl as a safener and in-built adjuvant system

**For selective grass weed control in autumn and  
spring sown barley, wheat, perennial ryegrass,  
pasture, and turf**

**NET CONTENTS: 10 LITRES**

©Registered Trademark of Arxada NZ Limited

Approved pursuant to the HSNO Act 1996, No HSR100112. See [www.epa.govt.nz](http://www.epa.govt.nz) for approval conditions.

Registered pursuant to the ACVM Act 1997, No. P009986 See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions.

**arxada**

	<b>GHS CLASSIFICATION</b>	<b>HAZARD STATEMENT</b>
	Skin sensitisation Category 1 Asp. Tox. 1 Specific target organ toxicity (repeated exposure) Category 2 Hazardous to the aquatic environment acute & chronic Cat. 1 Hazardous to Soil Organisms.	May cause an allergic skin reaction. May be fatal if swallowed and enters airways. May cause damage to organs through prolonged or repeated exposure. Very toxic to aquatic life with long lasting effects.

**FIRST AID:** IF SWALLOWED: Immediately call a POISON CENTER or doctor/physician. Do NOT induce vomiting.  
 IF ON SKIN: Wash with plenty of water. If skin irritation or rash occurs: Get medical advice/attention. Take off contaminated clothing and wash it before reuse.  
**PERSONAL PROTECTION:** DO NOT eat, drink or smoke while using. When preparing spray wear cotton overalls buttoned to the neck and wrist and washable hat, elbow-length PVC gloves and face shield or goggles. When using the prepared spray wear cotton overalls buttoned to the neck and wrist and washable hat and elbow-length PVC gloves. Remove protective clothing and wash hands and face thoroughly before meals and after work. Contaminated work clothing should not be allowed out of the workplace. Wash contaminated clothing before reuse. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.  
**STORAGE:** Store locked up. Store in original container, tightly closed and in a dry, cool area away from foodstuffs. Store in accordance with NZS 8409 Management of Agrichemicals. When stored appropriately this product should show no significant degradation for 2 years from the date of manufacture. Contact your supplier for further information about the use of this product if it is older than this.  
**DISPOSAL:** Avoid contamination of any water supply with product or empty container. Triple rinse empty container placing rinse water in the spray tank. If recycling, discard cap and deliver clean container to an Agrecovery depot or crush and bury in an approved landfill. Dispose of product only by using according to this label, or at an approved landfill or disposal facility.  
**SPILLAGE:** Wear appropriate protective clothing. Exclude all bystanders from the vicinity of the spill and prevent material from entering waterways. Contain liquid spills and absorb with inert material and place in waste container. Wash area with water and absorb with further inert material. Dispose of waste safely in an approved landfill.  
**SAFETY DATA SHEET:** Additional information is listed in the Safety Data Sheet available from Arxada NZ Limited or www.arxada.co.nz  
**QUALIFICATION REQUIREMENTS:** Persons mixing, loading, applying, or otherwise handling Jive must meet qualification requirements set out in the EPA Hazardous Substances (Hazardous Property Controls) Notice 2017.

**PLEASE READ THE ENTIRE LABEL BEFORE MIXING OR USE. SHAKE OR STIR WELL BEFORE USE.**

**GENERAL INFORMATION:**

**WEEDS CONTROLLED:** Wild oats, lesser canary grass (*Phalaris minor*), barnyard grass, rough bristle grass, summer grass, yellow bristle grass.  
**RESISTANT WEEDS:** Gnawed canary grass (*Phalaris paradoxa*), annual and hybrid ryegrasses, tall fescue, chewing's fescue, cocksfoot, browntop, grazing brome (=spiky brome – *Bromus stamineus*), prairie grass, and most broadleaved weeds are not controlled by Jive at label rates.

CROP	WEEDS	RATE	REMARKS
Barley, Wheat	Wild oats	750 mL/ha in 250 litres water	Apply when wild oats are between the 3 leaf and 1st node visible growth stage and when they are growing actively.
	Lesser canary grass ( <i>Phalaris minor</i> )		Apply when weeds are between the 3 leaf visible stage and the end of tillering and are growing actively. Avoid application during periods of moisture stress.
	Summer grass, Barnyard grass, Rough bristle grass		Apply to the seedling stage in spring before weeds have become reproductive. Control may be reduced if applications are made after stem elongation has begun.
Perennial ryegrass	Wild oats	750 mL/ha in 250 litres water	Apply in early spring when wild oats are between 3 leaf and end of tillering growth stage and growing actively. Note: Seed yield response is likely to be reduced the longer the delay in the removal of wild oats in spring. <i>Transient yellowing and suppression may occur in some ryegrass varieties. This may lead to a delay in heading however, final yield and biomass is unaffected.</i>
Pasture	Yellow bristle grass, Rough bristle grass, Summer grass, Barnyard grass	750 mL/ha in 250 litres water	Apply to established pasture 7 days after grazing during the period December/January. Apply to the seedling stage before grass weeds have become reproductive. Control will be reduced if applications are made after stem elongation.
Turf (ryegrass)	Yellow bristle grass, Rough bristle grass, Summer grass, Barnyard grass	Use 750 mL/ha in 250 litres of water as a broadcast spray or 15ml/ litre as a spot spray	Apply to established turf during spring and summer when conditions are warm and weeds are actively growing. Apply to the weed seedling stage before grass weeds have become reproductive.

**ADJUVANT:** Jive contains an inbuilt surfactant system. **DO NOT add any further surfactant or spraying oil.**  
**Ryegrass varieties tested:** A range of Perennial ryegrasses have been tested for crop safety. Please contact your local Arxada representative for full details on varieties tested.  
**DO NOT apply over Timothy (*Phleum pratense*)** as severe suppression may occur.

**TURF Species:**

**DO NOT** apply over seedling **Kikuyu** turf as complete plant loss may occur. Established kikuyu will be severely suppressed and partial or severe die back may occur.  
**DO NOT** apply over seedling **Paspalum** turf as severe suppression and complete plant loss may occur. Established Paspalum turf may be severely suppressed.

**MIXING:** Shake well before use. Add the required amount to a partly filled spray tank with agitation system operating. Complete filling the tank with water and apply.

**COMPATIBILITY:** DO NOT USE JIVE IN ANY MIXTURES WHEN APPLYING TO RYEGRASS SEEDCROPS. When used on barley and wheat, Jive is compatible with plant growth regulators. Care must be taken to ensure that recommended timings for all products are followed. The addition of adjuvant oil is NOT required. DO NOT use Jive within 5 days of broad-leaved herbicide applications.

**APPLICATION:** Do not overlap spray swathes in ryegrass crops as some minor crop suppression may occur in certain varieties. Flush spray system with clean water after use.

**RAINFASTNESS:** Jive is rain fast within one hour of application

**WITHHOLDING PERIODS:**

**It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.**

Barley and Wheat: 70 days (Do not graze treated crop for at least 70 days after treatment)

Perennial ryegrass: 42 days (Do not graze treated crop for at least 42 days after treatment)

Pasture 28 days (Do not graze treated pasture for at least 28 days after treatment)

**SOIL ACTIVITY:** Jive is rapidly broken down by soil microbes causing no restrictions on crop rotations.

**UNDERSOWN CROPS:** Jive can be used on crops undersown with certain grasses (See under "Resistant Weeds" above) and clover.

**RESISTANCE MANAGEMENT:** **GROUP 1 HERBICIDE** Jive contains fenoxaprop-P-ethyl, a selective grass herbicide of the aryloxyphenoxy-propionate group. Jive is a herbicide which inhibits production of acetyl CoA carboxylase. For weed resistance management Jive is a Group 1 herbicide. Some naturally occurring weed biotypes resistant to Jive, and other Group 1 herbicides, may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Jive or other Group 1 herbicides. See www.resistance.nzpps.org for more information.

**NOTICE TO BUYER:** The manufacturer and vendors of Jive warrant the formulation to contain the specified active ingredient within accepted tolerance but make no warranty of any kind on the uses of material and accept no responsibility, whether the material is used in accordance with the label directions or not.

**REGISTERED TO:** Arxada NZ Limited, Hudson Rd, New Plymouth, New Zealand.

Tel: 06 755-9234 Fax: 06 755-1174

**EMERGENCY PHONE: 0800 CHEMCALL 0800 243 622**

D.O.M:  
BATCH NO:

